

Inflammatory cytokines IL-1 β and TNF- α regulate p75^{NTR} expression in CNS neurons and astrocytes by distinct cell-type-specific signalling mechanisms

Soyoung Choi and Wilma J Friedman¹

Department of Biological Sciences, Rutgers University, Newark, NJ 07102, U.S.A.

Cite this article as: Choi S and Friedman WJ (2009) Inflammatory cytokines IL-1 β and TNF- α regulate p75^{NTR} expression in CNS neurons and astrocytes by distinct cell-type-specific signalling mechanisms. ASN NEURO 1(2):art:e00010.doi:10.1042/AN20090009

SUPPLEMENTARY ONLINE DATA




Figure S1 Confirmation that inhibitors of p38 MAPK and NF- κ B inhibit their respective pathways

Treatment of hippocampal astrocytes with (A) IL-1 β or (B) TNF- α in the absence or presence of the p38 MAPK inhibitor showed that SB203580 inhibited p38 MAPK phosphorylation induced by either ligand. (C) To confirm the inhibitory effect of SN-50 on NF- κ B nuclear translocation, astrocytes were pretreated with vehicle or with 10 μ M SN-50 for 30 min and then treated with IL-1 β or TNF- α for 30 min. Cells were fixed and stained for p65, and nuclei were labelled with Hoechst stain. Nuclear staining of p65 was observed in IL-1 β - or TNF- α -treated astrocytes without SN-50 (indicated by arrows; note the pink colour of the nuclei), and nuclear translocation of p65 was blocked by SN-50 (bottom panels).

¹To whom correspondence should be addressed (email wilmaf@andromeda.rutgers.edu).

© 2009 The Author(s) This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial Licence (<http://creativecommons.org/licenses/by-nc/2.5/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.




Figure S2 Hippocampal neurons were treated with or without biotin

Cells were processed as indicated in Figure 6 of the main paper, lysates were precipitated with streptavidin and analysed by Western blot for p75^{NTR} levels. In the absence of biotin, no p75^{NTR} was detected.

Received 2 February 2009/7 April 2009; accepted 22 April 2009

Published as Immediate Publication 22 April 2009, doi 10.1042/AN20090009